

7.4 Trigonometric Integrals

1. Let $u = \cos x$. Then $du = -\sin x dx$.

$$\begin{aligned}\int \cos^4 x \sin x dx &= -\int u^4 du \\ &= -\frac{u^5}{5} + C \\ &= -\frac{\cos^5 x}{5} + C\end{aligned}$$

2. Let $u = 5\phi$. Then $du = 5d\phi$.

$$\begin{aligned}\int \sin^2 5\phi d\phi &= \frac{1}{5} \int \sin^2 u du \\ &= \frac{1}{5} \int \frac{1}{2}(1 - \cos 2u) du \\ &= \frac{1}{10} \left(u - \frac{1}{2} \sin 2u \right) + C \\ &= \frac{1}{10} \left(5\phi - \frac{1}{2} \sin(2 \times 5\phi) \right) + C \\ &= \frac{1}{2} \phi - \frac{1}{20} \sin(10\phi) + C\end{aligned}$$

3. Let $u = \sin 2z$. Then $du = 2 \cos 2z dz$.

$$\begin{aligned}\int \sin^2 2z \cos^3 2z dz &= \int \sin^2 2z \cos^2 2z (1 - \sin^2 2z) dz \\ &= \int \sin^2 2z \cos 2z dz - \int \sin^4 2z \cos 2z dz \\ &= \frac{1}{2} \int u^2 du - \frac{1}{2} \int u^4 du \\ &= \frac{u^3}{6} - \frac{u^5}{10} + C \\ &= \frac{\sin^3 2z}{6} - \frac{\sin^5 2z}{10} + C\end{aligned}$$

4.

$$\begin{aligned}\int \sin x \cos\left(\frac{x}{2}\right) dx &= \int 2 \sin\left(\frac{x}{2}\right) \cos\left(\frac{x}{2}\right) \cos\left(\frac{x}{2}\right) dx \\ &= \int 2 \cos^2\left(\frac{x}{2}\right) \sin\left(\frac{x}{2}\right) dx\end{aligned}$$

Let $u = \cos\left(\frac{x}{2}\right)$. Then $du = -\frac{1}{2} \sin\left(\frac{x}{2}\right) dx$.

$$\begin{aligned}
 \int 2 \cos^2\left(\frac{x}{2}\right) \sin\left(\frac{x}{2}\right) dx &= -4 \int u^2 du \\
 &= -\frac{4u^3}{3} + C \\
 &= -\frac{4}{3} \cos^3\left(\frac{x}{2}\right) + C
 \end{aligned}$$

5. Let $u = \sec x$. Then $du = \sec x \tan x dx$.

$$\begin{aligned}
 \int \sec^4 x \tan^3 x dx &= \int \tan^2 x \sec^3 x (\sec x \tan x) dx \\
 &= \int (\sec^2 x - 1) \sec^3 x (\sec x \tan x) dx \\
 &= \int \sec^5 x (\sec x \tan x) dx - \int \sec^3 x (\sec x \tan x) dx \\
 &= \int u^5 du - \int u^3 du \\
 &= \frac{u^6}{6} - \frac{u^4}{4} + C \\
 &= \frac{\sec^6 x}{6} - \frac{\sec^4 x}{4} + C
 \end{aligned}$$

6.

$$\begin{aligned}
 \int \tan^4 x \sec x dx &= \int (\sec^2 x - 1)(\sec^2 x - 1) \sec x dx \\
 &= \int (\sec^5 x - 2 \sec^3 x + \sec x) dx \\
 &= \frac{\sec^3 x \tan x}{4} + \frac{3}{4} \int \sec^3 x dx - 2 \frac{\sec x \tan x}{2} - 2 \times \frac{1}{2} \int \sec x dx + \ln|\sec x + \tan x| + C \\
 &= \frac{\sec^3 x \tan x}{4} + \frac{3}{4} \left(\frac{\sec x \tan x}{2} + \frac{1}{2} \int \sec x dx \right) - 2 \frac{\sec x \tan x}{2} - \ln|\sec x + \tan x| + -\ln|\sec x + \tan x| + C \\
 &= \frac{\sec^3 x \tan x}{4} - \frac{5 \sec x \tan x}{8} + \frac{3}{8} \ln|\sec x + \tan x|
 \end{aligned}$$

7.

$$\begin{aligned}
 \int \sqrt{\tan x} \sec^4 x dx &= \int \tan^{\frac{1}{2}} x \sec^4 x dx \\
 &= \int \tan^{\frac{1}{2}} x (\tan^2 x + 1) \sec^2 x dx \\
 &= \int \left[\tan^{\frac{5}{2}} x \sec^2 x + \tan^{\frac{1}{2}} x \sec^2 x \right] dx
 \end{aligned}$$


Let $u = \tan x$. Then $du = \sec^2 x dx$.

$$\begin{aligned}
 \int \left[\tan^{\frac{5}{2}} x \sec^2 x + \tan^{\frac{1}{2}} x \sec^2 x \right] dx &= \int u^{\frac{5}{2}} du + \int u^{\frac{1}{2}} du \\
 &= \frac{u^{\frac{7}{2}}}{\frac{7}{2}} + \frac{u^{\frac{3}{2}}}{\frac{3}{2}} + C \\
 &= \frac{2 \tan^{\frac{7}{2}} x}{7} + \frac{2 \tan^{\frac{3}{2}} x}{3} + C
 \end{aligned}$$

8. Let $u = \frac{x}{2}$. Then $du = \frac{1}{2} dx$. If $x = 0$, then $u = 0$. If $x = \frac{\pi}{2}$, then $u = \frac{\pi}{4}$.

$$\begin{aligned}
 \int_0^{\frac{\pi}{2}} \tan^5 \left(\frac{x}{2} \right) dx &= \int_0^{\frac{\pi}{4}} 2 \tan^5 u \, du \\
 &= \frac{2 \tan^4 u}{4} \Big|_0^{\frac{\pi}{4}} - 2 \int_0^{\frac{\pi}{4}} \tan^3 u \, du \\
 &= 2 \left(\frac{\sqrt{2}}{2} \right)^4 - \frac{2 \tan^4 u}{2} \Big|_0^{\frac{\pi}{4}} + 2 \int_0^{\frac{\pi}{2}} \tan u \, du \\
 &= \frac{8}{16} - \left(\frac{\sqrt{2}}{2} \right)^2 - 2 \ln |\cos u| \Big|_0^{\frac{\pi}{4}} \\
 &= \frac{8}{16} - 1 - 2 \ln \left| \cos \left(\frac{\pi}{4} \right) \right| - 0 \\
 &= -\frac{1}{2} - 2 \ln \left| \frac{\sqrt{2}}{2} \right| \\
 &= -\frac{1}{2} + \ln \left| \frac{4}{2} \right| \\
 &= -\frac{1}{2} + \ln 2
 \end{aligned}$$

9.



$$\begin{aligned}
 V &= \pi \left(\int_0^{\frac{\pi}{4}} \cos^2 x \, dx - \int_0^{\frac{\pi}{4}} \sin^2 x \, dx \right) \\
 &= \pi \left[\frac{1}{2} \cos x \sin x + \frac{x}{2} \right]_0^{\frac{\pi}{4}} - \pi \left[-\frac{1}{2} \cos x \sin x + \frac{x}{2} \right]_0^{\frac{\pi}{4}} \\
 &= \pi \left[\left(\frac{1}{4} + \frac{\pi}{8} \right) - \left(-\frac{1}{4} + \frac{\pi}{8} \right) \right] \\
 &= \pi \left(\frac{1}{2} \right) \\
 &= \frac{\pi}{2}
 \end{aligned}$$

10.

a. $\int \csc x \, dx = \int \csc x \frac{\csc x + \cot x}{\csc x + \cot x} dx$

Let $u = \csc x + \cot x$. Then $du = (-\csc x \cot x - \csc^2 x) dx$.

$$\begin{aligned}
 \int \csc x \, dx &= - \int \frac{1}{u} du \\
 &= -\ln|u| + C \\
 &= -\ln|\csc x + \cot x| + C
 \end{aligned}$$

b.

$$\begin{aligned}
 \int \csc x \, dx &= \int \frac{1}{\sin x} dx \\
 &= \int \frac{\sin x}{\sin^2 x} dx \\
 &= \int \frac{\sin x}{(1 - \cos x)(1 + \cos x)} dx
 \end{aligned}$$

Let $u = \cos x$. Then $du = -\sin x \, dx$.

$$\begin{aligned}
\int \frac{\sin x}{(1 - \cos x)(1 + \cos x)} dx &= - \int \frac{du}{(1 - u)(1 + u)} \\
&= \int \frac{A}{1 - u} du + \int \frac{B}{1 + u} du \\
&= - \int \frac{\frac{1}{2}}{1 - u} du - \int \frac{\frac{1}{2}}{1 + u} du \\
&= \int \frac{\frac{1}{2}}{1 - \cos x} du - \int \frac{\frac{1}{2}}{1 + \cos x} du \\
&= \frac{1}{2} |1 - \cos x| - \frac{1}{2} |1 + \cos x| + C \\
&= \frac{1}{2} \left| \frac{1 - \cos x}{1 + \cos x} \right| + C \\
&= \ln \sqrt{\frac{1 - \cos x}{1 + \cos x}} + C \\
&= \ln \left| \tan \left(\frac{x}{2} \right) \right| + C
\end{aligned}$$

OR

$$\begin{aligned}
\ln \sqrt{\frac{1 - \cos x}{1 + \cos x}} + C &= \ln \sqrt{\frac{(1 - \cos x)}{(1 + \cos x)} \cdot \frac{(1 - \cos x)}{(1 - \cos x)}} + C \\
&= \ln \sqrt{\frac{(1 - \cos x)^2}{1 - \cos^2 x}} + C \\
&= \ln \left| \frac{1 - \cos x}{\sin x} \right| + C \\
&= \ln \left| \frac{1}{\sin x} - \frac{\cos x}{\sin x} \right| + C \\
&= \ln |\csc x - \cot x| + C
\end{aligned}$$